

Research

Open Access

Prolastin, a pharmaceutical preparation of purified human α 1-antitrypsin, blocks endotoxin-mediated cytokine release

Izabela Nita¹, Camilla Hollander², Ulla Westin² and Sabina-Marija Janciauskiene*¹

Address: ¹Department of Medicine, Lund University, University Hospital Malmö, 20502 Malmö, Sweden and ²Department of Otolaryngology and Head and Neck Surgery, Lund University, University Hospital Malmö, 20502 Malmö, Sweden

Email: Izabela Nita - izabela-nita@swipnet.se; Camilla Hollander - Camilla.Hollander@oron.mas.lu.se; Ulla Westin - Ulla.Peterson-Westin@oron.mas.lu.se; Sabina-Marija Janciauskiene* - sabina.janciauskiene@medforsk.mas.lu.se

* Corresponding author

Published: 31 January 2005

Received: 05 November 2004

Respiratory Research 2005, **6**:12 doi:10.1186/1465-9921-6-12

Accepted: 31 January 2005

This article is available from: <http://respiratory-research.com/content/6/1/12>

© 2005 Nita et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: α 1-antitrypsin (AAT) serves primarily as an inhibitor of the elastin degrading proteases, neutrophil elastase and proteinase 3. There is ample clinical evidence that inherited severe AAT deficiency predisposes to chronic obstructive pulmonary disease. Augmentation therapy for AAT deficiency has been available for many years, but to date no sufficient data exist to demonstrate its efficacy. There is increasing evidence that AAT is able to exert effects other than protease inhibition. We investigated whether Prolastin, a preparation of purified pooled human AAT used for augmentation therapy, exhibits anti-bacterial effects.

Methods: Human monocytes and neutrophils were isolated from buffy coats or whole peripheral blood by the Ficoll-Hypaque procedure. Cells were stimulated with lipopolysaccharide (LPS) or zymosan, either alone or in combination with Prolastin, native AAT or polymerised AAT for 18 h, and analysed to determine the release of TNF α , IL-1 β and IL-8. At 2-week intervals, seven subjects were submitted to a nasal challenge with sterile saline, LPS (25 μ g) and LPS-Prolastin combination. The concentration of IL-8 was analysed in nasal lavages performed before, and 2, 6 and 24 h after the challenge.

Results: *In vitro*, Prolastin showed a concentration-dependent (0.5 to 16 mg/ml) inhibition of endotoxin-stimulated TNF α and IL-1 β release from monocytes and IL-8 release from neutrophils. At 8 and 16 mg/ml the inhibitory effects of Prolastin appeared to be maximal for neutrophil IL-8 release (5.3-fold, $p < 0.001$ compared to zymosan treated cells) and monocyte TNF α and IL-1 β release (10.7- and 7.3-fold, $p < 0.001$, respectively, compared to LPS treated cells). Furthermore, Prolastin (2.5 mg per nostril) significantly inhibited nasal IL-8 release in response to pure LPS challenge.

Conclusion: Our data demonstrate for the first time that Prolastin inhibits bacterial endotoxin-induced pro-inflammatory responses *in vitro* and *in vivo*, and provide scientific bases to explore new Prolastin-based therapies for individuals with inherited AAT deficiency, but also for other clinical conditions.

Background

α 1-antitrypsin (AAT) is a glycoprotein, which is the major inhibitor of neutrophil elastase and proteinase 3 [1,2]. AAT is mainly produced in liver cells, but also in extrahepatic cells, such as monocytes, macrophages and pulmonary alveolar cells [3,4]. The average concentration of AAT in plasma in healthy individuals is 1.3 mg/ml, with a half-life of 3 to 5 days. AAT is an acute phase protein, and its circulating levels increase rapidly to concentrations exceeding 2 mg/ml in response to inflammation or infection [5]. Individuals with plasma AAT values below 0.7 mg/ml are considered to be AAT deficient [6,7]. Over 75 alleles of AAT have been identified to date, of which at least 20 affect either the amount or the function of the AAT molecule *in vivo* [6-8]. A very common deficiency allele is termed Z, which differs from the normal M in the substitution of Glu 342 to Lys [7,9,10]. This single amino acid exchange causes spontaneous polymerization of the AAT, markedly impeding its release into the circulation [11]. The retained material is associated with hepatic diseases [12], while diminished circulating levels lead to antiproteinase deficiency and higher susceptibility to elastase mediated tissue injury [13,14]. The alleles of AAT are inherited in an autosomal codominant manner [2]. Therefore, individuals heterozygous for the Z allele (MZ) have 30–40% whereas individuals homozygous for the Z allele (ZZ) have only 10–15% of normal plasma AAT levels [15-17]. Tobacco smoke and air pollution have long been recognised as risk factors for the development of chronic obstructive pulmonary disease (COPD); the only proven genetic risk factor, however, is the severe Z deficiency of AAT [18,19]. Cigarette smokers with AAT-deficiency develop COPD much earlier in life than smokers with the normal AAT genotype [8,10,11].

The pulmonary emphysema that is associated with inherited AAT deficiency is intimately linked with the lack of proteinase inhibitor within the lungs that is available to bind to, and inactivate, neutrophil elastase. On the basis of clinical observations involving patients with inherited AAT deficiency and various experimental studies, the elastase-AAT imbalance hypothesis became widely accepted as the explanation for lung tissue destruction in emphysema [20,21]. There is now increasing evidence that an excessive activity of various proteolytic enzymes in the lung milieu, including members of the serine, cysteine and metalloprotease families, may damage the elastin network of lungs [14]. Since the severe ZZ and intermediate MZ AAT deficiency accounts for less than 1–2% and 8–18% of emphysema cases, it is believed that the protease-antiprotease hypothesis provides a rational basis for the explanation of the development and progression of emphysema in general [22,23].

Based on the protease-antiprotease hypothesis, augmentation therapy of emphysema with severe AAT deficiency was introduced during the 1980s [24]. Intravenous administration of a pasteurized pooled human plasma AAT product (Prolastin; Bayer Corporation; Clayton, North Carolina) is used to increase AAT levels in deficient individuals [25]. The major concept behind augmentation therapy is that a rise in the levels of blood and tissue AAT will protect lungs from continuous destruction by proteases, particularly neutrophil elastase [26]. For example, anti-elastase capacity in the lung epithelial lining fluid has been found to increase to 60–70% of normal in homozygous Z AAT-deficient individuals subjected to augmentation therapy [26,27]. Whether this biochemical normalization of AAT levels influences the pathogenic processes of lung disease is still under debate. The most recent results, however, suggest that Prolastin therapy may have beneficial effects in reducing the frequency of lung infections and reducing the rate of decline of lung function [28,29].

There is growing evidence that AAT, in addition to its anti-proteinase activity, may have other functional activities. For example, AAT has been demonstrated to stimulate fibroblast proliferation and procollagen synthesis [30], to up-regulate human B cell differentiation into IgE- and IgG4-secreting cells [31], to interact with the proteolytic cascade of enzymes involved in apoptosis [32,33] and to express contrasting effects on the post-transcriptional regulation of iron between erythroid and monocytic cells [34]. AAT is also known to inhibit neutrophil superoxide production [35], induce macrophage-derived interleukin-1 receptor antagonist release [36] and reduce bacterial endotoxin and TNF α -induced lethality *in vivo* [37,38]. We recently demonstrated, *in vitro*, that both native (inhibitory) and non-inhibitory (polymerised and oxidised) forms of AAT strongly inhibit lipopolysaccharide-induced human monocyte activation [39]. AAT appears to act not just as an anti-proteinase, but as a molecule with broader anti-inflammatory properties. Data presented in this study provide clear evidence that Prolastin, a preparation used for AAT deficiency augmentation therapy, significantly inhibits bacterial endotoxin-induced pro-inflammatory cell responses *in vitro*, and suppresses nasal IL-8 release in lipopolysaccharide-challenged individuals, *in vivo*.

Materials and Methods

α 1-antitrypsin (AAT) preparations

α 1-antitrypsin (Human) Prolastin[®] (Lot 26N3PT2) was a gift from Bayer (Bayer Corporation, Clayton, North Carolina, USA). This vial of Prolastin contained 1059 mg of functionally active AAT, as determined by capacity to inhibit porcine pancreatic elastase. Prolastin was dissolved in sterile water for injections provided by

manufacture and stored at +4 °C. Purified human AAT was obtained from the Department of Clinical Chemistry, Malmö University Hospital, Sweden. Native AAT was diluted in phosphate buffered saline (PBS), pH 7.4. To ensure the removal of endotoxins, AAT was subjected to Detoxi-Gel AffinityPak columns according to instructions from the manufacturer (Pierce, IL, USA). Purified batches of AAT were then tested for endotoxin contamination with the Limulus amoebocyte lysate endochrome kit (Charles River Endosafe, SC, USA). Endotoxin levels were less than 0.2 enzyme units/mg protein in all preparations used. The concentrations of AAT in the endotoxin-purified batches were determined according to the Lowry method [40]. Polymeric AAT was produced by incubation at 60 °C for 10 h. Polymers were confirmed on non-denaturing 7.5% PAGE gels.

Monocyte isolation and culture

Monocytes were isolated from buffy coats using Ficoll-Paque PLUS (Pharmacia, Sweden). Briefly, buffy coats were diluted 1:2 in PBS with addition of 10 mM EDTA and layered on Ficoll. After centrifugation at 400 g for 35 min, at room temperature, the cells in the interface were collected and washed 3 times in PBS-EDTA. The cell purity and amount were determined in a cell counter Auto-counter AC900EO (Swelabs Instruments AB, Sweden). The granulocyte fractions were less than 10%. Cells were seeded into 12-well cell culture plates (Nunc, Denmark) at a concentration of 4×10^6 cells/ml in RPMI 1640 medium supplemented with penicillin 100 U/ml; streptomycin 100 µg/ml; non-essential amino acids 1×; sodium pyruvate 2 mM and HEPES 20 mM (Gibco, UK). After 1 h 15 min, non-adherent cells were removed by washing 3 times with PBS supplemented with calcium and magnesium. Fresh medium was added and cells were stimulated with lipopolysaccharide (LPS, 10 ng/ml, J5 Rc mutant; Sigma, Sweden) in the presence or absence of various concentrations of Prolastin (0–16 mg/ml), constant concentration of native or polymerised AAT (0.5 mg/ml) for 18 h at 37 °C, 5% CO₂.

Neutrophil isolation and culture

Human neutrophils were isolated from the peripheral blood of healthy volunteers using Polymorphprep TM (Axis-Shield PoC AS, Oslo, Norway) as recommended by the manufacture. In brief, 25 ml of anti-coagulated blood was gently layered over the 12.5 ml of Polymorphprep TM and centrifuged at 1600 rpm for 35 min. Neutrophils were harvested as a low band of the sample/medium interface, washed with PBS, and residual erythrocytes were subjected to hypotonic lysis. Purified neutrophils were washed in RPMI-1640- Glutamax-1 medium (Gibco-BRL Life Technologies, Grand Island, NY) supplemented with 0.1% bovine serum albumin (BSA) and resuspended in the same medium. The neutrophil purity was more than

75% as determined on an AutoCounter AC900EO. Cell viability was > 95% according to trypan blue staining.

Neutrophils (5×10^6 cells/ml) were plated into sterile endorph tubes. Zymosan was boiled, washed and sonicated. Opsonized zymosan was prepared by incubating zymosan with serum (1:3) in 37 °C water bath for 20 min. After, zymosan was centrifuged, washed with PBS and resuspended at 30 mg/ml. Cells alone or activated with zymosan (0.3 mg/ml) were exposed to various concentrations of Prolastin (0–8 mg/ml), and native or polymerised AAT preparations (0.5 mg/ml) for 18 h at 37 °C 5% CO₂. Cell free supernatants were obtained by centrifugation at 300 g for 10 min, and stored at -80 °C until analysis

Cytokine/chemokine analysis

Cell culture supernatants from monocytes and neutrophils stimulated with LPS or zymosan alone or in combination with Prolastin, native or polymerised AAT were analysed to determine TNFα, IL-1β and IL-8 levels by using DuoSet ELISA sets (R&D Systems, MN, USA; detection levels 15.6, 3.9, and 31.2 pg/ml, respectively).

Subjects

Seven subjects (four females and three males) of 26–50 (median 38) years of age, non-smokers, non-allergic volunteers participated in the study. All subjects gave written informed consent before participation in the study. None of the subjects has a history of respiratory disease and none took any medication at the study time.

Study Design

At 2-week intervals each subject was submitted to a nasal challenge with sterile saline, LPS and LPS-Prolastin combination. All experimental sessions were done in the same room. On each provocation day, the nose was inspected and cleaned with 8 ml of isotonic NaCl. Between nasal lavages the subjects stayed in the same building and asked to keep away from known sources of nasal irritants. The night was spent in their own homes. All participants completed a symptom questionnaire. In the first session, the baseline lavage was taken after instillation to each nostril of 8 ml of sterile isotonic NaCl. In the next session, the subjects were challenged with LPS from *Escherichia coli* serotype 026:B6, Lot 17H4042 (Sigma-Aldrich, USA). The provocation solution was prepared prior to use. LPS was added to 8 ml of sterile 0.9% NaCl to obtain a final concentration of 250 µg/ml, and 100 µl of the provocation solution was sprayed into each nostril, using a needle-less syringe. In the third session, the subjects were first challenged with LPS, as described above, and after 30 min with 2.5 mg of Prolastin into each nostril. Lavage samples were taken with instillation to each nostril of 8 ml of sterile isotonic NaCl after 2, 6 and 24 h followed by assessment of symptoms by a questionnaire. All subject

completed a symptom questionnaire with questions about nasal and eye irritation, and throat and airway symptoms. None of the participants reported symptoms of nasal, eye or throat irritations, and no general symptoms such as muscle pain, shivering, were mentioned.

Nasal Lavage

The procedure for nasal lavage was performed according to a method described by Wihl and co-workers [41]. Each nasal cavity was lavaged separately with a syringe (60 ml) to which a plastic nasal olive was connected for close nostril fitting. To prevent lavage spilling into the throat, the subject was bent forward at an angle of 60° during the procedure. Equilibrium was maintained between the mucosal lining and the lavage fluid by injecting the saline gently into the nasal cavity and drawing it back five times into the syringe. The lavage was performed in both nostrils and samples were collected into a test tube. The samples were then centrifuged at 1750 rpm, 6°C for 10 min and immediately frozen at -80°C. The protein concentration in the lavage fluids was measured by Lowry method and IL-8 levels were determined by DuoSet ELISA sets (R&D Systems, MN, USA; detection levels 31.2 pg/ml).

Statistical Analysis

Statistical Package (SPSS for Windows, release 11.5, SPSS Inc., Chicago) was used for the statistical calculations. The differences in the means of cell culture experimental results were analysed for their statistical significance with the one-way ANOVA combined with a multiple-comparisons procedure (Scheffe multiple range test). The equality of means of experimental results in healthy volunteers were analysed for statistical significance with independent two sample t-test and repeated measures of ANOVA using the SPSS MANOVA procedure <http://www.utexas.edu/cc/docs/stat38.html>. Tests showing $p < 0.05$ were considered to be significant.

Results

Concentration-dependent effects of Prolastin on LPS-induced cytokine release from human monocytes

Various concentrations of Prolastin (0–16 mg/ml) were added to adherent-isolated human monocytes with or without LPS (10 ng/ml). Cells stimulated with LPS alone served as a positive control, while PBS stimulated monocytes served as negative controls. As illustrated in figures 1A and 1B, simultaneous incubation of monocytes with LPS and Prolastin resulted in a reduction in TNF α and IL-1 β release compared to the cells stimulated with LPS alone. Inhibition of LPS-induced cytokine release by Prolastin was concentration-dependent and was typically observed over a concentration range of 0.5–16 mg/ml. At 16 mg/ml the inhibitory effects of Prolastin appeared to be maximal for both TNF α (10.7-fold, $p < 0.001$) and IL-1 β (7.3-fold, $p < 0.001$), compared to LPS alone.

Inhibitory effects at 0.5 mg/ml of AATs on LPS-mediated IL-1 β and TNF α release

We recently found that simultaneous incubation of monocytes with LPS and either the inhibitory (native) or non inhibitory (polymeric) form of AAT resulted in a reduction in TNF α and IL-1 β release compared to the cells stimulated with LPS alone. At 0.5 mg/ml the effects of native and polymerised AAT appeared to be maximal (41). Therefore, we selected a 0.5 mg/ml concentration of Prolastin, native and polymerised AAT, and compared their effects on LPS-stimulated cytokine release at 18 h. As shown in figures 2A and 2B, LPS triggered a significant release of TNF α and IL-1 β ($p < 0.001$ v medium alone) by monocytes. At 0.5 mg/ml, native and polymerised AAT remarkably inhibited LPS-induced TNF α and IL-1 β release ($p < 0.001$) (Fig. 2). The inhibitory effect of Prolastin (0.5 mg/ml) on LPS-stimulated TNF α release was comparable in magnitude to that of native or polymeric AAT, whereas its inhibitory effect on LPS-induced IL-1 β release did not reach significance.

Concentration-dependent effects of Prolastin on neutrophil IL-8 release


The effects of Prolastin (0–8 mg/ml) on human neutrophil IL-8 production are shown in Figure 3A. Neutrophils stimulated with opsonized zymosan (0.3 mg/ml) released a large amount of IL-8 ($p < 0.001$), compared to controls. Prolastin inhibited IL-8 release by neutrophils stimulated with opsonized zymosan (Fig 3A). This inhibition was concentration-dependant, with maximal suppression of IL-8 release (5.3-fold, $p < 0.001$ compared to zymosan treated cells) at 8 mg/ml.

Inhibitory effects at 0.5 mg/ml of native, polymeric AAT and Prolastin on zymosan-mediated IL-8 release


Neutrophils were stimulated with zymosan (0.3 mg/ml) or AATs (0.5 mg/ml) either alone or in combination for 18 h and IL-8 protein determined. As illustrated in figure 3B, polymeric and native AAT and Prolastin significantly inhibited the release of IL-8 protein by activated neutrophils. In terms of maximal effect, native AAT > polymerised AAT > Prolastin. It must be noted that native, polymeric AAT and Prolastin alone showed no effect on neutrophils, relative to non-treated buffer controls (data not shown).

Inhibition of the LPS-induced increase in nasal IL-8 release by Prolastin

To assess the effect of Prolastin on LPS-induced nasal provocation, IL-8 levels in nasal lavages were measured. Nasal instillation 25 μ g per nostril of LPS alone or in combination with 2.5 mg/ml of Prolastin was performed in non-smoking and non-allergic volunteers ($n = 7$, 4 females and 3 males). The IL-8 release in response to LPS challenge increased over time compared to baseline levels

**Figure 1**

A concentration-response inhibition of lipopolysaccharide-stimulated TNF α (A) and IL-1 β (B) release by Prolastin in human blood monocytes. Isolated blood monocytes were treated with LPS (10 ng/ml) alone or together with various concentrations of Prolastin (0–16 mg/ml) for 18 h. TNF α and IL-1 β levels were measured by ELISA. Data are the means of quadruplicate culture supernatants \pm S.E. and are representative of three separate experiments.

**Figure 2**

Comparisons of the effects of native (nAAT), polymeric (pAAT) and Prolastin on lipopolysaccharide – stimulated TNF α (A) and IL- β (B) production by human blood monocytes isolated from four healthy donors. Isolated blood monocytes were treated with LPS (10 ng/ml) alone or together with 0.5 mg/ml nAAT, pAAT or Prolastin for 18 h. TNF α and IL-1 β levels were measured by ELISA. Each bar represent the mean \pm S.E. *** $p < 0.001$.




Figure 3

Effects of AATs on neutrophils activated with zymosan. (A) Concentration-dependent effects of Prolastin on IL-8 release from neutrophils activated with opsonised zymosan. Freshly isolated blood neutrophils were treated with zymosan (0.3 mg/ml) alone or together with various concentrations of Prolastin (0–8 mg/ml) for 18 h. IL-8 levels were measured by ELISA. Data are the means of quadruplicate culture supernatants ± S.E. and are representative of three separate experiments. (B) Effects of opsonised zymosan alone or together with native (nAAT), polymeric (pAAT) AAT or Prolastin on IL-8 release from neutrophils. The release of neutrophil IL-8 was measured in cell free supernatants as described in Materials and methods. Neutrophils were treated for 18 h with a constant amount of zymosan (0.3 mg/ml) alone or together with nAAT, pAAT or Prolastin (0.5 mg/ml) for 18 h. IL-8 levels were measured by ELISA. Each bar represents the means ± S.E. of three separate experiments carried out in duplicate repeats. *** p < 0.001




Figure 4

IL-8 analysis in nasal lavage of subjects challenged with LPS alone or LPS+Prolastin combination. Seven healthy volunteers were treated with LPS (25 µg/nostril) or with LPS followed 30 min later with Prolastin (2.5 mg/nostril), nasal lavage was collected at different time points (0, 2, 6 and 24 h) as described in Material and Methods. The concentration of IL-8 (pg/ml) was measured by ELISA. IL-8 values are expressed as a ratio of IL-8 concentration at selected time point and the basal level. Independent two sample t-test shows after 6 and 24 h significantly higher levels of IL-8 in subjects treated with LPS compared to LPS+Prolastin. * $p < 0.05$

(Fig. 4). The levels of IL-8 increased already after 2 h of LPS challenge ($245.7\% \pm 87$) and remained higher after 24 h (310 ± 77.5) compared to baseline ($100\% \pm 19.2$). By contrast, when IL-8 levels were examined in LPS-Prolastin-treated lavage samples, no significant changes in IL-8 release were observed compared to baseline. In the presence of Prolastin, the LPS effect on IL-8 release was inhibited ($p < 0.05$) (Fig. 4).

Discussion

There is now, however, ample evidence that serine proteinase inhibitors (serpins), in addition to their well estab-

lished anti-inflammatory capacity to regulate serine proteinases activity, may possess broader anti-inflammatory properties. Several studies have shown that the biological responses of bacterial lipopolysaccharide (endotoxin) *in vivo* may be sensitive to serpins. For example, the serpin antithrombin, has been shown to protect animals from LPS-induced septic shock and also to inhibit IL-6 induction by LPS [42,43]. Our recent study provided first *in vitro* evidence that native (inhibitor) and at least two modified (non-inhibitory i.e. polymeric and oxidised) forms of AAT can block the release of an array of chemokine and cytokines from LPS-stimulated

monocytes [39]. These studies therefore further support a central role of serpins in inflammation, not only as the regulators of proteinase activity, but also as the suppressors of endotoxin induced pro-inflammatory responses. In line with these findings, we demonstrate here that Prolastin, a preparation of human AAT which is used for augmentation therapy, significantly inhibits endotoxin-induced pro-inflammatory effects *in vitro* and *in vivo*.

Stimulation of human monocytes and neutrophils with bacterial endotoxin results in the release of a range of inflammatory mediators including the pro-inflammatory cytokines (e.g. IL-6, IL-1 β and TNF α) and the chemokines (e.g. MCP-1 and IL-8) [44-46]. Together, these play a crucial role in the recruitment and activation of leukocytes and the subsequent release of harmful proteases that may further perpetuate the inflammatory process. We found that Prolastin significantly inhibits endotoxin-induced IL-1 β and TNF α release by monocytes and IL-8 release by neutrophils *in vitro*. The Prolastin exhibited these anti-inflammatory properties in a concentration-dependent manner. Its maximal effects were observed with 16 mg/ml in the monocyte model and with 8 mg/ml in the neutrophil model, since doubling these concentrations did not significantly modify the intensity of the effects. Indeed, Prolastin markedly prevented endotoxin-induced cell activation at 0.5–4 mg/ml concentrations, implying that these lower concentrations of Prolastin might also be sufficient to inhibit endotoxin effects. It is worth noting that in order to reduce a potential risk of transmission of infectious agents the Prolastin preparation is heat-treated in solution at 60° \pm 0.5 for not less than 10 h. Data from *in vitro* studies show that heat-treatment results in AAT polymerization and loss of its inhibitory activity [47,48]. Therefore, in our experimental model we compared anti-inflammatory effects of Prolastin with those of native and heat treated (60°C 10 h) AATs. At concentrations used (0.5 mg/ml), no significant difference was found between the effects of Prolastin and native or heat-treated (polymeric) AAT on endotoxin-induced monocyte TNF α and neutrophil IL-8 elevation. The median concentrations of endotoxin-stimulated IL-1 β levels also decreased in the presence of Prolastin but failed to reach statistical significance. In general, inhibitory effects on endotoxin-stimulated monocyte IL-1 β and neutrophil IL-8 release were better pronounced by native AAT compared to polymeric AAT or Prolastin. Similarly, in our previous study we found that in terms of maximal effect, native AAT > polymerised AAT > oxidized AAT were efficient in inhibiting LPS-stimulated TNF α and IL-1 β , and IL-8 release from monocytes [39]. Further studies will be necessary to better evaluate how temperature, pH or other physicochemical challenges may influence anti-inflammatory effectiveness of AAT preparations.

To explore our hypothesis that AAT functions as a potent inhibitor of endotoxin-induced effects, we examined whether Prolastin also inhibits responses to LPS in the nasal airway, *in vivo*. In particular, we were interested in concentrations of the neutrophil chemoattractant, IL-8. Endotoxin (or LPS) from gram-negative bacteria is a common air contaminant in a number of occupational conditions, especially those in which exposure to animal waste or plant matter occurs [44,49-51]. Levels of LPS in such environments may exceed 20 $\mu\text{g}/\text{m}^3$ air and may be associated with respiratory symptoms and nasal inflammation in exposed persons [52]. For example, nasal inflammation as evaluated by an increased influx of inflammatory cells into the nasal airway and increased IL-8 levels, has been described in persons occupationally exposed to LPS [51]. Moreover, it has been suggested that constitutive levels of IL-8 might further enhance responses to an inflammatory stimulus, such as LPS [53]. A number of experimental studies have shown that a nasal instillation of LPS causes the cytokine and chemokine reaction [54,55]. In our pilot study we also showed that instilled defined amounts of endotoxin (25 $\mu\text{g}/\text{per nostril}$) induce time-dependent nasal IL-8 release in normal subjects. Two hours after LPS instillation the IL-8 levels in nasal lavage reached more than twice the basal level and remained higher during all the times studied. However, during the next session, when 30 min after challenge with LPS, Prolastin (2.5 mg/ per nostril) was instilled, no induction of nasal IL-8 release was found compared to the basal levels. Furthermore, the protective ability of Prolastin did not disappear over study time. We cannot determine from these experiments whether Prolastin is directly suppressing IL-8 release or suppressing another inflammatory response that leads to IL-8 release; nonetheless, our finding suggests that effects of Prolastin directed against endotoxin-stimulated inflammatory responses may be beneficial.

Thus, data from both *in vitro* and *in vivo* experiments provide novel evidence that the Prolastin preparation is a potent inhibitor of endotoxin effects. The major concept behind augmentation therapy with pooled plasma-derived AAT has been that a rise in the level of AAT in subjects with severe inherited AAT deficiency would protect the lung tissue from continued destruction by proteinases (i.e. primarily leukocyte elastase) [7,56,57]. Recent findings provide evidence that augmentation therapy with AAT reduces the incidence of lung infections in patients with AAT-related emphysema [28,58]. Furthermore, Cantin and Woods have reported that aerosolized AAT suppresses bacterial proliferation in a rat model of chronic *Pseudomonas aeruginosa* lung infection [59]. Stockley and co-workers demonstrated that a short-term therapy of AAT augmentation not only restores airway concentrations of AAT to normal, but also reduces levels of leukotriene B₄, a major mediator of neutrophil recruitment and

activation. Interestingly, authors have suggested that the efficacy of AAT augmentation may be most beneficial in individuals with the most inflammation [29,60]. Data presented in this study clearly show that Prolastin inhibits endotoxin-stimulated pro-inflammatory responses, and thus provides new biochemical evidence supporting the efficacy of augmentation therapy. The current findings also suggest that Prolastin may, in fact, be used for broader clinical applications than merely augmentation therapy.

Abbreviations

AAT, α 1-antitrypsin; COPD, chronic obstructive pulmonary disease; LPS, lipopolysaccharide; ZZ, homozygous AAT-deficiency variant; MM, wild type AAT variant; PBS, phosphate buffered saline; EDTA, ethylenediaminetetraacetic acid; HEPES, 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid

Authors' contribution

Izabela Nita, performed cell culture experiments, made contribution to acquisition of data;

Camilla Hollander, made substantial contribution to patient study design, material collection and analysis; Ulla Westin, contributed to study design and data interpretation; Sabina Janciauskiene, contributed to conception and study design, data interpretation and wrote the article

Acknowledgments

This work was supported by grants from the Swedish Research Council, and Department of Medicine, Lund University, Sweden.

References

- Potempa J, Korzus E, Travis J: **The serpin superfamily of proteinase inhibitors: structure, function, and regulation.** *J Biol Chem* 1994, **269**:15957-15960.
- Brantly ML: **Alpha-1-antitrypsin genotypes and phenotypes.** In *Alpha-1-antitrypsin* Edited by: Crystal RG. New York, Marcel Dekker; 1996:45-59.
- Kalsheker N, Morley S, Morgan K: **Gene regulation of the serine proteinase inhibitors alpha-1-antitrypsin and alpha-1-antichymotrypsin.** *Biochem Soc Trans* 2002, **30**:93-98.
- Olsen GN, Harris JO, Castle JR, Waldman RH, Karmgard HJ: **Alpha-1-antitrypsin content in the serum, alveolar macrophages, and alveolar lavage fluid of smoking and nonsmoking normal subjects.** *J Clin Invest* 1975, **55**:427-430.
- Travis J, Shieh BH, Potempa J: **The functional role of acute phase plasma proteinase inhibitors.** *Tokai J Exp Clin Med* 1988, **13**:313-320.
- Crystal RG: **The alpha 1-antitrypsin gene and its deficiency states.** *Trends Genet* 1989, **5**:411-417.
- Hutchison DC: **Natural history of alpha-1-protease inhibitor deficiency.** *Am J Med* 1988, **84**:3-12.
- Needham M, Stockley RA: **Alpha 1-antitrypsin deficiency. 3: Clinical manifestations and natural history.** *Thorax* 2004, **59**:441-445.
- Lomas DA, Mahadeva R: **Alpha 1-antitrypsin polymerization and the serpinopathies: pathobiology and prospects for therapy.** *J Clin Invest* 2002, **110**:1585-1590.
- Luisetti M, Seersholm N: **Alpha-1-antitrypsin deficiency. 1: epidemiology of alpha-1-antitrypsin deficiency.** *Thorax* 2004, **59**:164-169.
- Carrell RW, Lomas DA: **Alpha 1-antitrypsin deficiency--a model for conformational diseases.** *N Engl J Med* 2002, **346**:45-53.
- Eriksson S: **Alpha 1-antitrypsin deficiency.** *J Hepatol* 1999, **30 Suppl 1**:34-39.
- Wiedemann HP, Stoller JK: **Lung disease due to alpha 1-antitrypsin deficiency.** *Curr Opin Pulm Med* 1996, **2**:155-160.
- Stockley RA: **Alpha-1-antitrypsin deficiency: what next?** *Thorax* 2000, **55**:614-618.
- Sandford AJ, Weir TD, Spinelli JJ, Pare PD: **Z and S mutations of the alpha-1-antitrypsin gene and the risk of chronic obstructive pulmonary disease.** *Am J Respir Cell Mol Biol* 1999, **20**:287-291.
- Talamo RC, Langley CE, Levine BV, Kazemi H: **Genetic vs. quantitative analysis of serum alpha 1-antitrypsin.** *N Engl J Med* 1972, **287**:1067-1069.
- Guenter CA, Welch MH, Ferguson S, Henderson L, Hammarsten JF: **Alpha-1-antitrypsin deficiency: heterozygosity, intermediate levels, and pulmonary disease.** *Chest* 1971, **59**:Suppl:16S+.
- Sandford AJ, Silverman EK: **Chronic obstructive pulmonary disease. 1: Susceptibility factors for COPD the genotype-environment interaction.** *Thorax* 2002, **57**:736-741.
- Chow CK: **Cigarette smoking and oxidative damage in the lung.** *Ann N Y Acad Sci* 1993, **686**:289-298.
- Laurell CB, Eriksson S: **The electrophoretic alpha-1-globulin pattern of serum in alpha-1-antitrypsin deficiency.** *Scand J Clin Lab Invest* 1963, **15**:132-140.
- Gross P, deTreville RT, Babyak MA, Kaschak M, Tolker EB: **Experimental emphysema: effect of chronic nitrogen dioxide exposure and of papain on normal and pneumoconiotic lungs.** *Aspen Emphysema Conf* 1967, **10**:357-378.
- Lieberman J, Winter B, Sastre A: **Alpha 1-antitrypsin Pi-types in 965 COPD patients.** *Chest* 1986, **89**:370-373.
- Lieberman J: **Intermediate antitrypsin deficiency.** *Am Rev Respir Dis* 1990, **141**:1078.
- Gadek JE, Klein HG, Holland PV, Crystal RG: **Replacement therapy of alpha 1-antitrypsin deficiency. Reversal of protease-antiprotease imbalance within the alveolar structures of PiZ subjects.** *J Clin Invest* 1981, **68**:1158-1165.
- Wewers MD, Casolaro MA, Sellers SE, Swayze SC, McPhaul KM, Wittes JT, Crystal RG: **Replacement therapy for alpha 1-antitrypsin deficiency associated with emphysema.** *N Engl J Med* 1987, **316**:1055-1062.
- Hubbard RC, Brantly ML, Sellers SE, Mitchell ME, Crystal RG: **Anti-neutrophil-elastase defenses of the lower respiratory tract in alpha 1-antitrypsin deficiency directly augmented with an aerosol of alpha 1-antitrypsin.** *Ann Intern Med* 1989, **111**:206-212.
- Stoller JK, Aboussouan LS: **alpha 1-Antitrypsin deficiency . 5: intravenous augmentation therapy: current understanding.** *Thorax* 2004, **59**:708-712.
- Lieberman J: **Augmentation therapy reduces frequency of lung infections in antitrypsin deficiency: a new hypothesis with supporting data.** *Chest* 2000, **118**:1480-1485.
- Stockley RA, Hill AT, Hill SL, Campbell EJ: **Bronchial inflammation: its relationship to colonizing microbial load and alpha(1)-antitrypsin deficiency.** *Chest* 2000, **117**:291S-3S.
- Dabbagh K, Laurent GJ, Shock A, Leoni P, Papakrivopoulou J, Chambers RC: **Alpha-1-antitrypsin stimulates fibroblast proliferation and procollagen production and activates classical MAP kinase signalling pathways.** *J Cell Physiol* 2001, **186**:73-81.
- Jeannin P, Lecoanet-Henchoz S, Delneste Y, Gauchat JF, Bonnefoy JY: **Alpha-1 antitrypsin up-regulates human B cell differentiation selectively into IgE- and IgG4- secreting cells.** *Eur J Immunol* 1998, **28**:1815-1822.
- Ikari Y, Mulvihill E, Schwartz SM: **alpha 1-Proteinase inhibitor, alpha 1-antichymotrypsin, and alpha 2-macroglobulin are the antiapoptotic factors of vascular smooth muscle cells.** *J Biol Chem* 2001, **276**:11798-11803.
- Daemen MA, Heemskerk VH, van't Veer C, Denecker G, Wolfs TG, Vandenabeele P, Buurman WA: **Functional protection by acute phase proteins alpha(1)-acid glycoprotein and alpha(1)-antitrypsin against ischemia/reperfusion injury by preventing apoptosis and inflammation.** *Circulation* 2000, **102**:1420-1426.
- Graziadei I, Gaggi S, Kaserbacher R, Braunsteiner H, Vogel W: **The acute-phase protein alpha 1-antitrypsin inhibits growth and proliferation of human early erythroid progenitor cells (burst-forming units-erythroid) and of human erythroleuke-**

- mic cells (K562) in vitro by interfering with transferrin iron uptake. *Blood* 1994, **83**:260-268.
35. Bucurenci N, Blake DR, Chidwick K, Winyard PG: **Inhibition of neutrophil superoxide production by human plasma alpha I-antitrypsin.** *FEBS Lett* 1992, **300**:21-24.
 36. Churg A, Dai J, Zay K, Karsan A, Hendricks R, Yee C, Martin R, MacKenzie R, Xie C, Zhang L, Shapiro S, Wright JL: **Alpha-I-antitrypsin and a broad spectrum metalloprotease inhibitor, RS113456, have similar acute anti-inflammatory effects.** *Lab Invest* 2001, **81**:1119-1131.
 37. Jie Z, Cai Y, Yang W, Jin M, Zhu W, Zhu C: **Protective effects of alpha I-antitrypsin on acute lung injury in rabbits induced by endotoxin.** *Chin Med J (Engl)* 2003, **116**:1678-1682.
 38. Libert C, Van Molle W, Brouckaert P, Fiers W: **alpha-I-Antitrypsin inhibits the lethal response to TNF in mice.** *J Immunol* 1996, **157**:5126-5129.
 39. Janciauskiene S, Larsson S, Larsson P, Virtala R, Jansson L, Stevens T: **Inhibition of lipopolysaccharide-mediated human monocyte activation, in vitro, by alpha I-antitrypsin.** *Biochem Biophys Res Commun* 2004, **321**:592-600.
 40. Lowry OH, Rosebrough NJ, Farr AL, Randall RJ: **Protein measurement with the Folin phenol reagent.** *J Biol Chem* 1951, **193**:265-275.
 41. Wihl JA, Baumgarten CR, Petersson G: **Contralateral differences among biomarkers determined by a modified nasal lavage technique after unilateral antigen challenge.** *Allergy* 1995, **50**:308-315.
 42. Dickneite G, Leithauser B: **Influence of antithrombin III on coagulation and inflammation in porcine septic shock.** *Arterioscler Thromb Vasc Biol* 1999, **19**:1566-1572.
 43. Souter PJ, Thomas S, Hubbard AR, Poole S, Romisch J, Gray E: **Antithrombin inhibits lipopolysaccharide-induced tissue factor and interleukin-6 production by mononuclear cells, human umbilical vein endothelial cells, and whole blood.** *Crit Care Med* 2001, **29**:134-139.
 44. Heumann D, Roger T: **Initial responses to endotoxins and Gram-negative bacteria.** *Clin Chim Acta* 2002, **323**:59-72.
 45. Harada A, Sekido N, Akahoshi T, Wada T, Mukaida N, Matsushima K: **Essential involvement of interleukin-8 (IL-8) in acute inflammation.** *J Leukoc Biol* 1994, **56**:559-564.
 46. Anderson P, Phillips K, Stoecklin G, Kederasha N: **Post-transcriptional regulation of proinflammatory proteins.** *J Leukoc Biol* 2004, **76**:42-47.
 47. Chow MK, Devlin GL, Bottomley SP: **Osmolytes as modulators of conformational changes in serpins.** *Biol Chem* 2001, **382**:1593-1599.
 48. Huntington JA, Pannu NS, Hazes B, Read RJ, Lomas DA, Carrell RW: **A 2.6 Å structure of a serpin polymer and implications for conformational disease.** *J Mol Biol* 1999, **293**:449-455.
 49. Schwartz DA, Thorne PS, Yagla SJ, Burmeister LF, Olenchock SA, Watt JL, Quinn TJ: **The role of endotoxin in grain dust-induced lung disease.** *Am J Respir Crit Care Med* 1995, **152**:603-608.
 50. Schwartz DA, Thorne PS, Jagielo PJ, White GE, Bleuer SA, Frees KL: **Endotoxin responsiveness and grain dust-induced inflammation in the lower respiratory tract.** *Am J Physiol* 1994, **267**:L609-17.
 51. Peden DB, Tucker K, Murphy P, Newlin-Clapp L, Boehlecke B, Hazucha M, Bromberg P, Reed W: **Eosinophil influx to the nasal airway after local, low-level LPS challenge in humans.** *J Allergy Clin Immunol* 1999, **104**:388-394.
 52. Blaski CA, Watt JL, Quinn TJ, Thorne PS, Schwartz DA: **Nasal lavage cellularity, grain dust, and airflow obstruction.** *Chest* 1996, **109**:1086-1092.
 53. Sigsgaard T, Bonefeld-Jorgensen EC, Kjaergaard SK, Mamas S, Pedersen OF: **Cytokine release from the nasal mucosa and whole blood after experimental exposures to organic dusts.** *Eur Respir J* 2000, **16**:140-145.
 54. Danuser B, Rebsamen H, Weber C, Krueger H: **Lipopolysaccharide-induced nasal cytokine response: a dose-response evaluation.** *Eur Arch Otorhinolaryngol* 2000, **257**:527-532.
 55. Besancon-Watelet C, Bene MC, Montagne P, Faure GC, Jankowski R: **Eosinophilia and cell activation mediators in nasal secretions.** *Laryngoscope* 2002, **112**:43-46.
 56. Stoller JK: **Alpha-I antitrypsin deficiency: an under-recognized but important issue for respiratory therapists.** *Respir Care* 2003, **48**:1022-1024.
 57. Hill AT, Bayley DL, Campbell EJ, Hill SL, Stockley RA: **Airways inflammation in chronic bronchitis: the effects of smoking and alpha I-antitrypsin deficiency.** *Eur Respir J* 2000, **15**:886-890.
 58. Dirksen A, Dijkman JH, Madsen F, Stoel B, Hutchison DC, Ulrik CS, Skovgaard LT, Kok-Jensen A, Rudolphus A, Seersholm N, Vrooman HA, Reiber JH, Hansen NC, Heckscher T, Viskum K, Stolk J: **A randomized clinical trial of alpha(1)-antitrypsin augmentation therapy.** *Am J Respir Crit Care Med* 1999, **160**:1468-1472.
 59. Cantin AM, Woods DE: **Aerosolized prolactin suppresses bacterial proliferation in a model of chronic Pseudomonas aeruginosa lung infection.** *Am J Respir Crit Care Med* 1999, **160**:1130-1135.
 60. Stockley RA, Bayley DL, Unsal I, Dowson LJ: **The effect of augmentation therapy on bronchial inflammation in alpha I-antitrypsin deficiency.** *Am J Respir Crit Care Med* 2002, **165**:1494-1498.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

