

RESEARCH

Open Access

L-citrulline supplementation reverses the impaired airway relaxation in neonatal rats exposed to hyperoxia

Ramadan B Sopi^{1,2*}, Syed IA Zaidi^{3,4*}, Mitko Mladenov², Hazbije Sahiti¹, Zahide Istrefi¹, Icko Gjorgoski², Azem Lajçi¹ and Muharrem Jakupaj¹

Abstract

Background: Hyperoxia is shown to impair airway relaxation via limiting L-arginine bioavailability to nitric oxide synthase (NOS) and reducing NO production as a consequence. L-arginine can also be synthesized by L-citrulline recycling. The role of L-citrulline supplementation was investigated in the reversing of hyperoxia-induced impaired relaxation of rat tracheal smooth muscle (TSM).

Methods: Electrical field stimulation (EFS, 2–20 V)-induced relaxation was measured under *in vitro* conditions in precontracted tracheal preparations obtained from 12 day old rat pups exposed to room air or hyperoxia (>95% oxygen) for 7 days supplemented with L-citrulline or saline (*in vitro* or *in vivo*). The role of the L-citrulline/L-arginine cycle under basal conditions was studied by incubation of preparations in the presence of argininosuccinate synthase (ASS) inhibitor [α -methyl-D, L-aspartate, 1 mM] or argininosuccinate lyase inhibitor (ASL) succinate (1 mM) and/or NOS inhibitor [N^{ω} -nitro-L-arginine methyl ester; 100 μ M] with respect to the presence or absence of L-citrulline (2 mM).

Results: Hyperoxia impaired the EFS-induced relaxation of TSM as compared to room air control ($p < 0.001$; $0.5 \pm 0.1\%$ at 2 V to $50.6 \pm 5.7\%$ at 20 V in hyperoxic group; 0.7 ± 0.2 at 2 V to $80.0 \pm 5.6\%$ at 20 V in room air group). Inhibition of ASS or ASL, and L-citrulline supplementation did not affect relaxation responses under basal conditions. However, inhibition of NOS significantly reduced relaxation responses ($p < 0.001$), which were restored to control level by L-citrulline. L-citrulline supplementation *in vivo* and *in vitro* also reversed the hyperoxia-impaired relaxation. The differences were significant ($p < 0.001$; $0.8 \pm 0.3\%$ at 2 V to $47.1 \pm 4.1\%$ at 20 V without L-citrulline; $0.9 \pm 0.3\%$ at 2 V to $68.2 \pm 4.8\%$ at 20 V with L-citrulline). Inhibition of ASS or ASL prevented this effect of L-citrulline.

Conclusion: The results indicate the presence of an L-citrulline/L-arginine cycle in the airways of rat pups. L-citrulline recycling does not play a major role under basal conditions in airways, but it has an important role under conditions of substrate limitations to NOS as a source of L-arginine, and L-citrulline supplementation reverses the impaired relaxation of airways under hyperoxic conditions.

Keywords: Airway relaxation, Airway smooth muscle, Argininosuccinate lyase, Argininosuccinate synthase, L-arginine, Nitric oxide, Nitric oxide synthase

* Correspondence: ramadan.sopi@uni-pr.edu; syed.zaidi@case.edu

¹Department of Pharmacy-Biology, Faculty of Medicine, University of Prishtina, St. Martyrs' Boulevard n.n., Prishtina 10000, Kosovo

³Department of Chemical Engineering, Case Western Reserve University, Cleveland, OH 44106, USA

Full list of author information is available at the end of the article

Background

Bronchopulmonary dysplasia (BPD) is a chronic respiratory disease that results from complications related to the lung injury during treatment of respiratory distress syndrome in premature babies. Increased airway reactivity in childhood is the most common long-term manifestation of neonatal lung injury and is pronounced when there is a history of BPD [1,2]. While a variety of pre- and postnatal inflammatory processes may contribute to its etiology, exposure of an immature respiratory tract to increased supplemental oxygen plays an important contributing role to the development of BPD [3]. Therefore, neonatal hyperoxic exposure may serve as a model for BPD in which enhanced airway reactivity is a well-known consequence. We and others have demonstrated in rat pups that exposure to high inspired oxygen is associated with many pathophysiological features of BPD, including increased contractile responses and decreased relaxant responses of airways under *in vitro* and *in vivo* conditions using trachea and lung parenchymal tissues, as well as changes in lung morphology [4-11].

Nitric oxide (NO) is a potent relaxant of airway smooth muscle (ASM) [12,13]. Exposure of rat pups to high oxygen concentration has been shown to impair relaxant responses of the tracheal smooth muscle (TSM) [8] and the distal airways using lung parenchymal tissue [4,5]. The decreased NO production was implicated in hyperoxia-induced impairment of relaxation of airways NO-cGMP-dependent mechanism [14]. In order to compensate for the decrease of endogenous NO in neonatal care, exogenous NO is frequently used to prevent the adverse effects of neonatal hyperoxia [14-16]. Recent randomized human trials suggest that inhaled NO (iNO) decreases the incidence of BPD in subsets of premature babies [3,17,18]. However, inconsistent efficacy [19], complexity of iNO delivery in nonintubated patients, and high cost provide rationales for efficacious alternatives to iNO.

As NO is produced from L-arginine via NO synthase (NOS) [20], bioavailability of L-arginine plays a key role in the production of NO and on lung function. In most cells, L-arginine requirements are met primarily by the uptake of extracellular L-arginine via specific cationic amino acid transporters (system γ^+) [21]. However, in NO-producing cells, L-arginine may also be obtained through the L-citrulline/L-arginine cycle [22]. Through this cycle, L-citrulline can be recycled as L-arginine to the NOS pathway by the successive action of argininosuccinate synthase (ASS), which converts L-citrulline and L-aspartate into L-argininosuccinate, and then the argininosuccinate lyase (ASL) hydrolyses L-argininosuccinate into L-arginine and fumarate [22-25], (Figure 1). Some authors suggest that L-citrulline might be an indirect precursor of NO in NO-synthesizing cells [26]. L-citrulline has been shown to restore impaired relaxation of airways after early asthmatic reaction [27]. Also in other tissues, including the opossum

internal anal sphincter and rat gastric fundus, it has been shown that the L-citrulline/L-arginine cycle is involved in inhibitory nonadrenergic-noncholinergic nerve (iNANC)-mediated smooth muscle relaxation, under conditions of limited L-arginine availability to nNOS, which is induced by competitive inhibitors of the enzyme [28-31]. Furthermore, it was demonstrated that L-citrulline prevents O₂-inducing alveolar damage and alters angiogenesis in developing lungs [11]. However, there are no studies showing the role of the L-citrulline/L-arginine cycle on hyperoxia-induced impairment of ASM relaxation. The study of L-citrulline becomes even more important in hyperoxic condition because, due to the increased arginase activity [32], conditions of limited L-arginine availability for NOS are created. Therefore, this study was undertaken to test the hypothesis that the L-citrulline/L-arginine cycle is active in rat pup airways and that L-citrulline supplementation will reverse hyperoxia-induced impaired relaxation of ASM.

Methods

Animals and tissue preparation

Sprague-Dawley rat pups were used for these experiments. On the 5th day of life (P5), weight-matched rat pups from different litters were randomly mixed and assigned to either hyperoxic ($n=32$) or room air ($n=43$) groups. We have shown earlier that hyperoxic exposure for seven days starting from day 5 shows BPD-like symptoms in rat pups [4]. Therefore, the pups were exposed to hyperoxia (>95% O₂) or room air for seven days. Hyperoxic animals were housed in a Plexiglas chamber and exposed to a continuous flow of O₂ (2 L/min), whereas control animals were kept in a commercial rat cage in room air. Mothers were rotated each day between room air and hyperoxic groups to protect them from the toxicity of a constant hyperoxic exposure. Oxygen concentration was monitored continuously via oxygen analyzer (Pro O₂ Nuvair, Oxnard, CA). In one set of experiments, the animals were injected intraperitoneally (i.p.) with 200 mg/kg/day L-citrulline during hyperoxic exposure. Control animals were injected with the same volume of saline. No differences in weight were observed between the compared groups.

Animals were euthanized by asphyxiation in CO₂ on day 12 of postnatal life (P12). The trachea was removed and prepared with the intact epithelium, free of serosal connective tissue, in an ice-cold oxygenated Krebs-Henseleit (KH) solution (concentration in mM: 118.2 NaCl, 25 NaHCO₃, 4.6 KCl, 1.2 KH₂PO₄, 1.2 MgSO₄, 2.5 CaCl₂, and 10% dextrose, pH=7.4). A cylindrical airway segment of 3-mm length was isolated from the mid-portion of the tracheas of each animal, and then suspended between a stainless hook at the bottom of the organ bath and a force displacement transducer (FT 03 C, Grass Instruments, Quincy, MA). TSM tension was measured by a Power Lab/4SP AD Instruments, monitored and recorded using




Chart 4.0 software. An initial load of 0.3 g was used, and then tissues were allowed to equilibrate for 40 min in the organ baths containing KH solution (20 ml) at 37°C. The solution was continuously aerosolized with a gas mixture of 95% O₂ and 5% CO₂. All protocols described in this study were approved by the Internal Animal Use and Care Committees of the University of Prishtina and Case Western Reserve University.

Electrical field stimulation-induced relaxation experiments

After equilibration, a cumulative concentration-response curve was made to find the concentration of bethanechol that elicited 50–75% of maximal response in TSM (100 μM bethanechol). Incremental electrical field stimulation (EFS) was applied for 10s at 2-min intervals to induce relaxation through platinum electrodes to precontracted TSM at various voltages (2–20 V, AC). The relaxation of cylinders after EFS was expressed as a percentage of the total precontracted state for each cylinder [4].

An NOS inhibitor – N^ω-nitro-L-arginine methyl ester (L-NAME; 100 μM) and L-citrulline (2 mM) were applied to the organ bath 30 min prior to the addition of bethanechol, while an ASS inhibitor–α-methyl-D,L-aspartate (α-MDLA;

1 mM) and an ASL inhibitor–succinate (1 mM) were applied 40 min before bethanechol.


Statistical analysis

The results are expressed as mean ± SEM. Statistical significance was determined by Two-way ANOVA with repeated measurements. To analyze the differences in treatments at different voltage settings between groups and within a group, a *post-hoc* Tukey–Kramer multiple comparison test was used. In all cases, $p < 0.05$ was considered statistically significant.

Results

Effect of hyperoxia on ASM relaxation

Consistent with our previous studies, EFS-induced relaxation increased with the increasing voltages and hyperoxia significantly reduced the EFS-induced relaxation of precontracted TSM ($n = 10$) as compared to room air ($p < 0.001$; $n = 10$). The relaxant responses in the hyperoxic group ranged from $0.5 \pm 0.1\%$ at 2 V to $50.6 \pm 5.7\%$ at 20 V while those in the control group ranged from $0.7 \pm 0.2\%$ at 2 V to $80.0 \pm 5.6\%$ at 20 V (Figure 2). The differences became significant at ≥ 12 V



(Figure 2). Overall there was a significant difference between incremental voltages within the groups ($p < 0.01$).

Role of L-citrulline, α -MDLA and succinate on EFS-induced TSM relaxation in room air-exposed animals

To study the role of the L-citrulline/L-arginine cycle under basal conditions, the tissues were incubated with L-citrulline (2 mM), or α -MDLA (1 mM), or succinate (1 mM), or in combination of L-citrulline with succinate. There was no effect of L-citrulline on the EFS-induced relaxation of precontracted TSM of rat pups when compared to control responses ($n = 7$ per group). The α -MDLA, succinate or combination of L-citrulline and succinate did not produce any significant effect on EFS-induced relaxant responses ($n = 7$ per group). The relaxation values of TSM obtained from control tissues ranged from $1.1 \pm 0.4\%$ at 2 V to $78.1 \pm 4.0\%$ at 20 V (Figure 3). These data indicate that the L-citrulline/L-arginine cycle does not play a major role in the airway relaxation of rat pups under basal conditions (Figure 3).

Effect of L-citrulline, α -MDLA, and succinate on TSM relaxation after NOS blockage in room air-exposed animals

To study the effect of L-citrulline recycling under conditions of reduced utilization of L-arginine by NOS, tracheal rings obtained from the room air-exposed animals ($n = 8$) were preincubated with an NOS inhibitor—L-NAME (100 μ M), and then constricted using bethanecol. L-NAME significantly reduced ($p < 0.01$) the EFS-induced relaxation of TSM compared to the control responses, particularly at higher voltages (8–20 V; $p < 0.01$), (Figure 4). The data ranged from $0.7 \pm 0.2\%$ at 2 V to $50.5 \pm 7.2\%$ at 20 V, whereas in the control responses, the data ranged from $0.5 \pm 0.2\%$ at 2 V to $78.0 \pm 4.9\%$ at 20 V. Incubation of tissues with L-citrulline (2 mM) reversed the inhibition of EFS-induced relaxation of TSM produced by L-NAME ($p < 0.01$), indicating a critical role of L-citrulline under limiting

conditions of NO production. These data ranged from $0.6 \pm 0.1\%$ at 2 V to $72.1 \pm 4.0\%$ at 20 V, (Figure 4). α -MDLA (1 mM) prevented the capability of L-citrulline to restore diminished relaxant responses of TSM due to L-NAME. These differences were significant ($p < 0.01$) when compared to responses obtained from tissues coincubated in L-NAME plus L-citrulline. The data ranged from $0.4 \pm 0.2\%$ at 2 V to $52.3 \pm 5.2\%$ at 20 V. Similarly, succinate significantly ($p < 0.01$) prevented the effect of L-citrulline to restore the reduced relaxant responses of TSM due to L-NAME. The data ranged from $0.8 \pm 0.4\%$ at 2 V to $57.9 \pm 4.7\%$ at 20 V (Figure 5).

Effect of L-citrulline, α -MDLA and succinate on TSM relaxation in hyperoxia-exposed rat pups

As shown above, hyperoxia impaired the EFS-induced relaxation of rat pup TSM ($p < 0.001$) as compared to room air-exposed animals ($n = 7$ per group). The data ranged from $0.8 \pm 0.3\%$ at 2 V to $47.1 \pm 4.1\%$ at 20 V in hyperoxic animals and $0.9 \pm 0.2\%$ at 2 V to $74.8 \pm 5.6\%$ at 20 V in room air animals. L-citrulline supplementation (2 mM) reversed the hyperoxia-induced impaired relaxation ($p < 0.001$). The data ranged $0.9 \pm 0.3\%$ at 2 V to $68.2 \pm 4.8\%$ at 20 V (in hyperoxia + L-citrulline group) The differences were significant at higher voltages ≥ 8 V ($p < 0.001$). Both the ASS inhibitor, α -MDLA (1 mM) and the ASL inhibitor, succinate (1 mM) prevented L-citrulline from reversing the hyperoxia-induced impairment of relaxant responses (Figure 6 and Figure 7). The data ranged from $0.6 \pm 0.1\%$ at 2 V to $53.1 \pm 3.5\%$ at 20 V (in the presence of α -MDLA) and from $0.7 \pm 0.2\%$ at 2 V to $47.9 \pm 4.1\%$ at 20 V (in the presence of succinate), respectively.

Intraperitoneal supplementation of L-citrulline reversed the hyperoxia-induced impaired relaxation

As shown in Figure 8 *i.p.* supplementation of rat pups with L-citrulline (200 mg/kg/day) reversed the impaired relaxation in hyperoxic animals to normal levels compared to hyperoxic controls ($p < 0.01$). The differences were significant at higher voltages (≥ 12 V).

Discussion

Prior research by us has shown that the long-term exposure of hyperoxia impairs airway relaxation [4,32] and enhances airway contraction in neonatal rat lung [5]. In this study we have shown that hyperoxia decreases EFS-induced relaxant responses of ASM. Both of these processes complement each other in increasing airway hyperreactivity. Hyperoxia increases arginase activity, which limits the availability of L-arginine to NOS [4]. Limitation of L-arginine, being a common substrate for arginase and NOS, disrupts the NO/cGMP signaling pathway, resulting in impaired relaxation [4,32]. In our recent publication we have shown that production of NO in ASM cells is reduced in hyperoxia-




Figure 3 Role of L-citrulline, α -MDLA, and succinate on EFS-induced relaxation of TSM of rat pups exposed to room air: No significant effect of L-citrulline, α -MDLA and L-succinate individually or in combination was observed in TSM of room air exposed rat pups ($n = 7$ for each protocol of the experiments).

exposed rat pups [32]. In a lung cell, L-arginine is used as a substrate for NOS and is obtained from different sources for the production of NO, including the recycling of L-citrulline [31]. Maarsingh *et al.* [27] have shown the existence of an L-citrulline/L-arginine cycle in TSM. We have observed that the supplementation of L-citrulline reversed the hyperoxia-induced impaired relaxation in both *in vivo* (i.p. administration to animals) and *in vitro* (addition to organ bath) conditions. The *in vitro* study indicates that in rat TSM, there is an active L-citrulline/L-arginine cycle that produces L-arginine. Since the systemic administration of L-citrulline via i.p. route is also effective, we assume that, unlike L-arginine, L-citrulline is not taken up by the liver from the portal circulation, but is metabolized in the kidney to L-arginine that increases the plasma concentration of L-arginine. For this reason, oral or systemic administration

of L-citrulline could be a better substitute for L-arginine as it bypasses the liver [22].

The inhibition of the enzymes (ASL and ASS) involved in the recycling pathway of L-citrulline to L-arginine did not affect EFS-induced relaxation of TSM under basal conditions, indicating that L-arginine in basal conditions is not a limiting substrate. Furthermore, the supplementation of TSM with exogenous L-citrulline did not affect EFS-induced relaxation, indicating that the supply of L-arginine for NOS through L-citrulline cycling is not an important source for NOS substrate under these conditions. However, this pathway appears to play a key role once the NOS enzyme is blocked. Interestingly, L-citrulline supplementation fully restored the reduced relaxation caused by the blockage of NOS under basal conditions. The effect of L-citrulline was more prominent particularly at higher voltages when an




Figure 4 Effect of L-citrulline and α -MDLA on TSM relaxation after NOS blockade: EFS-induced relaxation of TSM preparations obtained from room air-exposed rat pups was significantly ($p < 0.01$) reduced by the addition of $100 \mu\text{M}$ L-NAME. The decrease in relaxation due to L-NAME was returned to normal levels by the addition of L-citrulline (2 mM). The argininosuccinate synthase (ASS) inhibitor – α -MDLA (1 mM) prevented the capability of L-citrulline to recover the diminished relaxation responses due to L-NAME. * *Room air + L-NAME vs. Room air*; § *Room air + L-NAME + L-citrulline vs. Room air + L-NAME*; † *Room air + L-NAME + L-citrulline + α -MDLA vs. Room air + L-NAME + L-citrulline*. ** $p < 0.01$; § $p < 0.05$; §§ $p < 0.01$; † $p < 0.05$; †† $p < 0.01$. ($n = 8$ for each protocol of the experiments).




Figure 5 Effect of L-citrulline and succinate on TSM relaxation after NOS blockade: EFS-induced relaxation of TSM preparations obtained from room air-exposed rat pups was significantly ($p < 0.01$) reduced by the addition of 100 μ M L-NAME. The decrease in relaxation due to L-NAME was returned to normal levels by the addition of L-citrulline (2 mM). The argininosuccinate lyase (ASL) inhibitor – succinate (1 mM) prevented the capability of L-citrulline to recover the diminished relaxation responses due to L-NAME. * Room air + L-NAME vs. Room air; § Room air + L-NAME + L-citrulline vs. Room air + L-NAME; † Room air + L-NAME + L-citrulline + succinate vs. Room air + L-NAME + L-citrulline. ** $p < 0.01$; § $p < 0.05$; §§ $p < 0.01$; † $p < 0.05$; †† $p < 0.01$. ($n = 8$ for each protocol of the experiments).

extra supply of NO is required to compensate the changes. This observation suggests that L-citrulline competes with the NOS blocker (L-NAME) reversing its inhibitory effects. These findings are comparable with other studies performed in the tracheal rings of a guinea pig model of asthma and human ASM using single or multiple frequencies of EFS [3,27], where the inhibitory effect of NOS inhibitor, N^o-nitro-L-arginine, on NO-mediated iNANC relaxation was restored by L-citrulline as well as L-arginine [33]. In addition, the inhibition of ASL reversed this beneficial effect of L-citrulline supplementation, indicating the presence of an active L-citrulline/L-arginine cycle in the airways of rat pups under conditions of substrate limitation for NO production. Similar results were shown in the tracheal rings of guinea pigs [27].

In asthmatic patients, a decrease in plasma L-arginine levels and an increase in serum arginase activity was observed [34]. Similarly, in the experimental BPD in neonatal rats, the plasma level of L-arginine was found to have decreased after hyperoxic exposure of animals [11], while arginase protein expression was upregulated in the airway epithelium, further depriving the adjacent ASM for L-arginine substrate. Furthermore, Mallese *et al.* [35] showed that hyperoxia diminishes NO production in mice because of the increase in hepatic arginase activity. L-citrulline also plays a key role in the reversal of impaired relaxation during hyperoxia that creates a L-arginine limiting condition. L-citrulline has been reported to be a noncompetitive inhibitor of arginase [36]. Therefore, in hyperoxic conditions, L-citrulline appears to work through arginase




Figure 6 Effect of L-citrulline and inhibition of argininosuccinate synthase (ASS) on TSM relaxation in hyperoxia-exposed rat pups: L-citrulline supplementation (2 mM) reversed the hyperoxia-induced impairment of relaxation of TSM ($p < 0.001$). The effect of the L-citrulline was lost when TSM was incubated in the presence of argininosuccinate synthase (ASS) inhibitor, α -MDLA (1 mM) ($p < 0.05$). *Hyperoxia vs. Room air; § Hyperoxia + L-citrulline vs. Hyperoxia; †Hyperoxia + L-citrulline + α -MDLA vs. Hyperoxia + L-citrulline. * $p < 0.05$; *** $p < 0.001$; §§§ $p < 0.001$; † $p < 0.05$ ($n = 7$ for each protocol of the experiments).




Figure 7 Effect of L-citrulline and inhibition of argininosuccinate lyase (ASL) on TSM relaxation in hyperoxia-exposed rat pups: The addition of 2 mM L-citrulline reversed the hyperoxia-induced impairment of relaxation of TSM ($p < 0.001$). The effect of the L-citrulline was lost when TSM was incubated in the presence of argininosuccinate lyase (ASL) inhibitor, L-succinate (1 mM) ($p < 0.01$). *Hyperoxia vs. Room air; § Hyperoxia + L-citrulline vs. Hyperoxia; †Hyperoxia + L-citrulline + succinate vs. Hyperoxia + L-citrulline. * $p < 0.05$; *** $p < 0.001$; §§ $p < 0.01$; §§§ $p < 0.001$; † $p < 0.05$; †† $p < 0.01$ ($n = 7$ for each protocol of the experiments).

inhibition rather than competition with NOS inhibitors as discussed above.

In a rat pup model of BPD, L-arginine becomes a limiting substrate and disrupts the NO-cGMP signaling pathway, due to the hyperoxia-induced upregulation of arginase activity. The disruption of NO-cGMP signaling reduces the relaxation of airways as a consequence [4,32]. An increase of arginase activity has been linked to airway hyperreactivity under different conditions in different animal models, such as rat pups, mice, and guinea pigs, as well as in humans [34,35,37-39]. In the perfused trachea of guinea pigs, it was demonstrated that increased arginase activity contributes to the airway hyperresponsiveness after early asthmatic reaction and deficiency of cNOS-derived NO [40]. Marsingh *et al.* [41] have shown that the increase of arginase activity after early asthmatic reaction in guinea pigs impairs the neuronal nitric oxide-mediated relaxation of ASM. Therefore, in this study the role of L-citrulline recycling in the

airways under *in vitro* and *in vivo* conditions of TSM obtained from hyperoxia exposed animals was studied. Hyperoxia-induced impairment of the relaxant responses of TSM of rat pups was reversed by L-citrulline supplementation under both *in vitro* and *in vivo* conditions, indicating the role of L-citrulline/L-arginine recycling enzymes in hyperoxic conditions. In order to demonstrate that the recovery of the impaired relaxation of TSM was due to recycling of L-citrulline into L-arginine, the ASS and ASL were inhibited using specific inhibitors. Interestingly, the protective effect of L-citrulline in hyperoxic TSM was lost by the inhibition of ASS, as well as ASL activity, further confirming the role of the L-citrulline/L-arginine cycle and the substrate limitation to NOS. Maarsingh *et al.* [27] also have found that exogenous L-citrulline fully reversed the impaired iNANC relaxation in the guinea pig model of asthma. We have shown that after hyperoxic exposure, arginase activity increased in the experimental BPD model [4,32]. This model demonstrates that the




Figure 8 Effect of *in vivo* L-citrulline supplementation on TSM relaxation in hyperoxia-exposed rat pups: *In vivo* supplementation of rat pups with L-citrulline (200 mg/kg/day) reversed the hyperoxia-induced impairment of relaxant responses of TSM in the hyperoxic animals ($p < 0.01$). *Hyperoxia + L-citrulline vs. Hyperoxia; * $p < 0.05$; ** $p < 0.01$ ($n = 4$ for each protocol of the experiments).

raised levels of arginase activity, which compete with NOS for common substrate, contribute to the impairment of relaxation in most distal airways because of the decrease of NO production [4,32].

Since it has been shown that the limitation of L-arginine availability to NOS is involved in pathogenesis of several diseases involving NO deficiencies, L-arginine supplementation to the animals has been used to treat these disorders. We have observed that the supplementation of 300 mg/kg of L-arginine single dose given daily for seven days through the intraperitoneal route reversed the impaired relaxant responses in rat pups [32]. Although L-arginine supplementation in rat pups exposed to hyperoxia seems to normalize impaired relaxation of lung parenchymal strips, it further increased arginase activity, thus reducing the effectiveness of L-arginine therapy. In addition, oral L-arginine administration failed to reduce airway hyperresponsiveness in a murine model of allergic asthma [42] or in asthmatic patients [43]. This failure might be because of the presystemic elimination of L-arginine by increased intestinal or hepatic arginase activity [44].

In contrast, L-citrulline is not taken up by the liver, but is metabolized predominantly in the kidney to L-arginine and could be considered as a masked precursor of L-arginine bypassing the liver [24], thus providing plenty of substrate for NO-cGMP signaling pathway [45]. The benefits of oral administration of L-citrulline have been shown in many diseases. Oral administration of L-citrulline in patients suffering from sickle cell disease raised the decreased plasma levels of L-arginine without any toxicity [46]. Furthermore, L-citrulline decreased pulmonary hypertension after surgery in patients with congenital heart disease [47], and it also reduced blood pressure response to cold stress [48]. Recently, Cormio *et al.* [49] have shown that oral L-citrulline supplementation improves erection hardness in men with mild erectile dysfunction. Another study revealed that supplementation of L-citrulline prevents hyperoxia-induced lung injury in newborn rats [11]. L-citrulline also ameliorates development of pulmonary hypertension and increases NO production in piglets exposed to chronic hypoxia [50]. These studies provide evidence that support the use of L-citrulline as a therapy in diseases characterized by NO deficiency.

The animal model used in this study has proven quite successful to characterize the role of the L-citrulline/L-arginine cycle under normal and limiting conditions of NOS substrate. We recognize that this model under *in vitro* conditions has limitations, as it is isolated from systemic circulation and lacks the connections from central nervous system, which also affect the physiology of airways.

Conclusion

This study demonstrates that the prolonged hyperoxic exposure to new born rats causes impaired relaxation of tracheal smooth muscle. Supplementation of L-citrulline

in vitro and *in vivo* reverses hyperoxia-induced attenuated relaxant responses of rat pup TSM in L-arginine limiting conditions. However, L-citrulline has no effect on relaxation in basal conditions when L-arginine is not a limiting factor. The evidence originated from this study suggests that supplementation with L-citrulline may have higher potential than L-arginine as a therapeutic use for preterm infants at risk for increased airway reactivity.

Abbreviations

ASM: Airway smooth muscle; α -MDLA: α -methyl-D,L-aspartate; ASL: Argininosuccinate lyase; ASS: Argininosuccinate synthase; BPD: Bronchopulmonary dysplasia; EFS: Electrical field stimulation; iNANC: Inhibitory non-adrenergic non-cholinergic; KH: Krebs-Henseleit; L-NAME: N^o-nitro-L-arginine methyl ester; nNOS: Neuronal; NOS; NO: Nitric oxide; NOS: Nitric oxide synthase; TSM: Tracheal smooth muscle.

Competing interest

The authors do not have any competing interests.

Authors' contributions

RBS designed and coordinated the study, performed a major part of the experiments, and drafted the manuscript. SIAZ participated in designing the study, carried out a part of experiments, and contributed to finalizing the manuscript. MM substantially participated in experiments and revised the manuscript. HS performed the *in vivo* experiments. ZI assisted in the experiments, analyzed the data, supervised the statistics, and contributed to the manuscript. IG contributed to the study design, data interpretation, and final revision of the manuscript. AL advised during the experiments and contributed to finalizing the manuscript. MJ conceived the study and participated in its design and direction, as well as preparing the manuscript. All authors read and approved the final draft of the manuscript.

Acknowledgements

This study was supported partially by the Ministry of Health of the Republic of Kosovo [MSH-90300576/11] and an IKARIA grant to SIAZ.

Author details

¹Department of Pharmacy-Biology, Faculty of Medicine, University of Prishtina, St. Martyrs' Boulevard n.n., Prishtina 10000, Kosovo. ²Institute of Biology, Faculty of Natural Sciences and Mathematics, "Sts, Cyril and Methodius" University, Skopje 1000, Macedonia. ³Department of Chemical Engineering, Case Western Reserve University, Cleveland, OH 44106, USA. ⁴Department of Chemical Engineering, Case Western Reserve University, Cleveland, OH 44106, USA.

Received: 13 January 2012 Accepted: 27 July 2012

Published: 7 August 2012

References

1. Pelkonen AS, Hakulinen AL, Turpeinen M: **Bronchial lability and responsiveness in school children born very preterm.** *Am J Respir Crit Care Med* 1997, **156**:1178–1184.
2. Hack M, Taylor HG, Drotar D, Schluchter M, Cartar L, Andreias L, Wilson-Costello D, Klein N: **Chronic conditions, functional limitations and special health care needs of school-aged children born with extremely low-birth-weight in the 1990's.** *JAMA* 2005, **294**:318–325.
3. Kinsella JP, Greenough A, Abman SH: **Bronchopulmonary dysplasia.** *Lancet* 2006, **367**:1421–1431.
4. Sopi RB, Haxhiu MA, Martin RJ, Dreshaj IA, Kamath S, Zaidi SI: **Disruption of NO-cGMP signaling by neonatal hyperoxia impairs relaxation of lung parenchyma.** *Am J Physiol Lung Cell Mol Physiol* 2007, **293**:L1029–L1036.
5. Sopi RB, Martin RJ, Haxhiu MA, Dreshaj IA, Yao Q, Jafri A, Zaidi SI: **Role of brain-derived neurotrophic factor in hyperoxia-induced enhancement of contractility and impairment of relaxation in lung parenchyma.** *Am J Physiol Lung Cell Mol Physiol* 2008, **295**:L348–L355.
6. Belik J, Jankov RP, Pan J, Tanswell AK: **Chronic O₂ exposure enhances vascular and airway smooth muscle contraction in the newborn but not adult rat.** *J Appl Physiol* 2003, **94**:2303–2312.

7. Hershenson MB, Abe MK, Kelleher MD, Naureckas ET, Garland A, Zimmermann A, Rubinstein VJ, Solway J: **Recovery of airway structure and function after hyperoxic exposure in immature rats.** *Am J Respir Crit Care Med* 1994, **149**:1663–1669.
8. Mhanna MJ, Haxhiu MA, Jaber MA, Walenga RW, Chang CH, Liu S, Martin RJ: **Hyperoxia impairs airway relaxation in immature rats via a cyclic AMP-mediated mechanism.** *J Appl Physiol* 2004, **96**:1854–1860.
9. Iben SC, Dreshaj IA, Farver CF, Haxhiu MA, Martin RJ: **Role of endogenous nitric oxide in hyperoxia-induced airway hyperreactivity in maturing rats.** *J Appl Physiol* 2000, **89**:1205–1212.
10. Iben SC, Haxhiu MA, Farver CF, Miller MJ, Martin RJ: **Short-term mechanical ventilation increases airway reactivity in rat pups.** *Pediatr Res* 2006, **60**:136–140.
11. Vadivel A, Aschner JL, Rey-Parra GJ, Magarik J, Zeng H, Summar M, Eaton F, Thébaud B: **L-citrulline attenuates arrested alveolar growth and pulmonary hypertension in oxygen-induced lung injury in newborn rats.** *Pediatr Res* 2010, **68**:519–525.
12. Fisher JT, Haxhiu MA, Martin RJ: **Regulation of lower airway function.** In *Fetal and Neonatal Physiology*. 2nd edition. Edited by Polin RA, Fox WW. Orlando, FL: Saunders; 1998:1060–1070.
13. Sherman TS, Chen Z, Yuhanna IS, Lau KS, Margraf LR, Shaul PW: **Nitric oxide synthase isoform expression in the developing lung epithelium.** *Am J Physiol* 1999, **276**:L383–390.
14. Cotton RB, Sundell HW, Zeldin DC, Morrow JD, Roberts LJ, Hazinski TA, Law AB, Steele S: **Inhaled nitric oxide attenuates hyperoxic lung injury in lambs.** *Pediatr Res* 2006, **59**:142–146.
15. Auten RL, Mason SN, Whorton MH, Lampe WR, Foster WM, Goldberg RN, Li B, Stamler JS, Auten KM: **Inhaled ethyl nitrite prevents hyperoxia-impaired postnatal alveolar development in newborn rats.** *Am J Respir Crit Care Med* 2007, **176**:291–299.
16. Rose MJ, Stenger MR, Joshi MS, Welty SE, Bauer JA, Nelin LD: **Inhaled nitric oxide decreases leukocyte trafficking in the neonatal mouse lung during exposure to >95% oxygen.** *Pediatr Res* 2010, **67**:244–9.
17. Ballard RA, Truog WE, Canan A, Martin RJ, Ballard PL, Merrill JD, Walsh MC, Durand DJ, Mayock DE, Eichenwald EC, Nuli DR, Hudak ML, Puri AR, Golombek SG, Courtney SE, Stewart DL, Welty SE, Phibbs RH, Hibbs AM, Luan X, Wadlinger SR, Asselin JM, Coburn CE: **Inhaled nitric oxide in preterm infants undergoing mechanical ventilation.** *N Engl J Med* 2006, **355**:343–353.
18. Schreiber MD, Gin-Mestran K, Marks JD, Huo D, Lee G, Srisuparp P: **Inhaled nitric oxide in premature infants with the respiratory distress syndrome.** *N Engl J Med* 2003, **349**:2099–2107.
19. Cole SF, Alleyne C, Barks JDE, Boyle RJ, Carroll JL, Dokken D, Edwards WH, Georgieff M, Gregory K, Johnston MV, Kramer M, Mitchell C, Neu J, De Pursley WM, Robinson WM, Rowitch DH: **NIH Consensus Development Conference Statement: Inhaled Nitric-Oxide Therapy for Premature Infants.** *Pediatrics* 2011, **127**:363.
20. Murad F: **Shattuck Lecture: Nitric oxide and cyclic GMP in cell signaling and drug development.** *N Engl J Med* 2006, **355**:2003–2011.
21. Closs EI, Simon A, Vekony N, Rotmann A: **Plasma membrane transporters for arginine.** *J Nutr* 2004, **134**:2752S–2759S.
22. Morris SM: **Regulation of enzymes of the urea cycle and arginine metabolism.** *Annu Rev Nutr* 2002, **22**:87–105.
23. Wiesinger H: **Arginine metabolism and the synthesis of nitric oxide in the nervous system.** *Prog Neurobiol* 2001, **64**:365–391.
24. Curis E, Nicolis I, Moinard C, Osowska S, Zerrouk N, Benazeth S, Cynober L: **Almost all about citrulline in mammals.** *Amino Acids* 2005, **29**:177–205.
25. Husson A, Brasse-Lagnel C, Fairand A, Renouf S, Lavoinne A: **Argininosuccinate synthetase from the urea cycle to the citrulline-NO cycle.** *Eur J Biochem* 2003, **270**:1887–1899.
26. Mori M, Gotoh T: **Regulation of nitric oxide production by arginine metabolic enzymes.** *Biochem Biophys Res Commun* 2000, **275**:715–719.
27. Maarsingh H, Leusink J, Zaagsma J, Meurs H: **Role of the L-citrulline/L-arginine cycle in iNANC nerve-mediated nitric oxide production and airway smooth muscle relaxation in allergic asthma.** *Eur J Pharmacol* 2006, **546**:171–176.
28. Chakder S, Rattan S: **L-arginine deficiency causes suppression of nonadrenergic noncholinergic nerve-mediated smooth muscle relaxation: role of L-citrulline recycling.** *J Pharmacol Exp Ther* 1997, **282**:378–384.
29. Rattan S, Chakder S: **L-citrulline recycling in opossum internal anal sphincter relaxation by nonadrenergic, noncholinergic nerve stimulation.** *Gastroenterology* 1997, **112**:1250–1259.
30. van Geldre LA, Timmermans JP, Lefebvre RA: **L-citrulline recycling by argininosuccinate synthetase and lyase in rat gastric fundus.** *Eur J Pharmacol* 2002, **455**:149–160.
31. Wu G, Morris SM Jr: **Arginine metabolism: nitric oxide and beyond.** *Biochem J* 1998, **336**:1–17.
32. Ali NK, Jafri A, Sopi RB, Prakash YS, Martin RJ, Zaidi SI: **Role of Arginase in Impairing Relaxation of Lung Parenchyma of Hyperoxia-Exposed Neonatal Rats.** *Neonatology* 2011, **101**:106–115.
33. Ellis JL, Conanan N: **L-citrulline reverses the inhibition of nonadrenergic, noncholinergic relaxations produced by nitric oxide synthase inhibitors in guinea pig trachea and human bronchus.** *J Pharmacol Exp Ther* 1994, **269**:1073–1078.
34. Morris CR, Poljakovic M, Lavrisha L, Machado L, Kuypers FA, Morris SM: **Decreased arginine bioavailability and increased serum arginase activity in asthma.** *Am J Respir Crit Care Med* 2004, **170**:148–153.
35. Mallek DT, Rogers LK, Velluci SM, Young TL, Park MS, Long DW, Welty SE, Smith CV, Nelin LD: **Hyperoxia increases hepatic arginase expression and ornithine production in mice.** *Toxicol Appl Pharmacol* 2006, **215**:109–117.
36. Shearer JD, Richards JR, Mills CD, Caldwell MD: **Differential regulation of macrophage arginine metabolism: A proposed role in wound healing.** *Am J Physiol* 1997, **272**:E181–190.
37. Que LG, Kantrow SP, Jenkinson CP, Piatadosi CA, Huang YCT: **Induction of arginase isoforms in the lung during hyperoxia.** *Am J Physiol Lung Cell Mol Physiol* 1998, **275**:L96–L102.
38. Meurs H, Maarsingh H, Zaagsma J: **Arginase and asthma: novel insights into nitric oxide homeostasis and airway hyperresponsiveness.** *Trends Pharmacol Sci* 2003, **24**:450–455.
39. Maarsingh H, Tio MA, Zaagsma J, Meurs H: **Arginase attenuates inhibitory nonadrenergic noncholinergic nerve-induced nitric oxide generation and airway smooth muscle relaxation.** *Respi Res* 2005, **6**:23.
40. Meurs H, McKay S, Maarsingh H, Hamer MA, Macic L, Molendijk N, Zaagsma J: **Increased arginase activity underlies allergen-induced deficiency of cNOS-derived nitric oxide and airway hyperresponsiveness.** *Br J Pharmacol* 2002, **136**:391–398.
41. Maarsingh H, Leusink J, Bos IS, Zaagsma J, Meurs H: **Arginase strongly impairs neuronal nitric oxide-mediated airway smooth muscle relaxation in allergic asthma.** *Respi Res* 2006, **7**:6.
42. Takano H, Lim HB, Miyabara Y, Ichinose T, Yoshikawa T, Sagai M: **Oral administration of L-arginine potentiates allergen-induced airway inflammation and expression of interleukin-5 in mice.** *J Pharmacol Exp Ther* 1998, **286**:767–771.
43. de Gouw HW, Verbruggen MB, Twiss IM, Sterk PJ: **Effect of oral L-arginine on airway hyperresponsiveness to histamine in asthma.** *Thorax* 1999, **54**:1033–1035.
44. Morris SM Jr: **Enzymes of arginine metabolism.** *J Nutr* 2004, **134**:2743S–2747S.
45. Schwedhelm E, Maas R, Freese R, Jung D, Lukacs Z, Jambrechina A, Spickler W, Schulze F, Böger RH: **Pharmacokinetic and pharmacodynamic properties of oral L-citrulline and L-arginine: impact on nitric oxide metabolism.** *Br J Clin Pharmacol* 2008, **65**:51–59.
46. Waugh WH, Daeschner CW, Files BA, McConnell ME, Strandjord SE: **Oral citrulline as arginine precursor may be beneficial in sickle cell disease: early phase two results.** *J Natl Med Assoc* 2001, **93**:363–371.
47. Smith HA, Canter JA, Christian KG, Drinkwater DC, Scholl FG, Christman BW, Rice GD, Barr FE, Summar M: **Nitric oxide precursors and congenital heart surgery: a randomized controlled trial of oral citrulline.** *J Thorac Cardiovasc Surg* 2006, **132**:58–65.
48. Figueroa A, Trivino JA, Sanchez-Gonzalez MA, Vicil F: **Oral L-citrulline supplementation attenuates blood pressure response to cold pressor test in young men.** *Am J Hypertens* 2010, **23**:12–16.
49. Cormio L, De Siati M, Lorusso F, Selvaggio O, Mirabella L, Sanguedolce F, Carrieri G: **Oral L-citrulline supplementation improves erection hardness in men with mild erectile dysfunction.** *Urology* 2011, **77**:119–122.
50. Ananthkrishnan M, Barr FE, Summar ML, Smith HA, Kaplowitz M, Cunningham G, Magarik J, Zhang Y, Fike CD: **L-Citrulline ameliorates chronic hypoxia-induced pulmonary hypertension in newborn piglets.** *Am J Physiol Lung Cell Mol Physiol* 2009, **297**:L506–L511.

doi:10.1186/1465-9921-13-68

Cite this article as: Sopi et al.: L-citrulline supplementation reverses the impaired airway relaxation in neonatal rats exposed to hyperoxia. *Respiratory Research* 2012 **13**:68.